



P-ISSN: 2349-8528

E-ISSN: 2321-4902

IJCS 2019; 7(1): 667-670

© 2019 IJCS

Received: 21-11-2018

Accepted: 25-12-2018

#### Kalaiyaranan C

Department of Agronomy,  
Faculty of Agriculture,  
Annamalai University, New  
Delhi, India

#### J Jayaraman

Department of Agronomy,  
Faculty of Agriculture,  
Annamalai University, New  
Delhi, India

#### S Jawahar

Department of Agronomy,  
Faculty of Agriculture,  
Annamalai University, New  
Delhi, India

#### MV Sriramachandrasekharan

Department of Soil science and  
Agricultural chemistry,  
Faculty of Agriculture,  
Annamalai University, New  
Delhi, India

#### Suseendran K

Department of Agronomy,  
Faculty of Agriculture,  
Annamalai University, New  
Delhi, India

#### S Elankavi

Department of Agronomy,  
Faculty of Agriculture,  
Annamalai University, New  
Delhi, India

#### Correspondence

#### Kalaiyaranan C

Department of Agronomy,  
Faculty of Agriculture,  
Annamalai University, New  
Delhi, India

## International Journal of Chemical Studies

# Integrated Sulphur management on yield, quality, nutrient uptake and nutrient availability of sunflower

**Kalaiyaranan C, J Jayaraman, S Jawahar, MV Sriramachandrasekharan, Suseendran K and S Elankavi**

### Abstract

Field investigations were carried out during March 2012 and August 2012 at Experimental Farm, Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalai Nagar, to study the effect of Integrated sulphur management practices on the yield, quality, nutrient uptake and nutrient availability of hybrid sunflower cv. sunbred. The experiments were laid out in randomized block design and replicated thrice. The experiment consisted of eleven treatments viz., T<sub>1</sub> – No sulphur / RDF alone (control), T<sub>2</sub> – 45 kg S ha<sup>-1</sup> through gypsum, T<sub>3</sub> – 45 kg S ha<sup>-1</sup> through vermicompost, T<sub>4</sub> – 45 kg S ha<sup>-1</sup> through poultry manure, T<sub>5</sub> – 45 kg S ha<sup>-1</sup> through lignite flyash, T<sub>6</sub> – 75 % S ha<sup>-1</sup> through gypsum + 25 % S ha<sup>-1</sup> through vermicompost, T<sub>7</sub> – 75 % S ha<sup>-1</sup> through gypsum + 25 % S ha<sup>-1</sup> through poultry manure, T<sub>8</sub> – 75 % S ha<sup>-1</sup> through gypsum + 25 % S ha<sup>-1</sup> through lignite flyash, T<sub>9</sub> – 50 % S ha<sup>-1</sup> through gypsum + 50 % S ha<sup>-1</sup> through vermicompost, T<sub>10</sub> – 50 % S ha<sup>-1</sup> through gypsum + 50 % S ha<sup>-1</sup> through poultry manure and T<sub>11</sub> – 50 % S ha<sup>-1</sup> through gypsum + 50 % S ha<sup>-1</sup> through lignite flyash. The results of the experiments revealed that application of 75 % S ha<sup>-1</sup> through gypsum + 25 % S ha<sup>-1</sup> through vermicompost (T<sub>6</sub>) significantly influenced the yield, quality, nutrient uptake and nutrient availability in sunflower. This treatment also recorded higher values for yield attributes viz., total number of seeds head<sup>-1</sup> (778.06 and 815.26), number of filled deeds head<sup>-1</sup> (634.45 and 672.28), seed filling percent (83.04 and 84.26 %), seed yield (1825.46 and 1927.39 kg ha<sup>-1</sup>) and stalk yield (3800.63 and 3898.59 kg ha<sup>-1</sup>) in both the crops. This treatment recorded maximum N,P,K and S uptake, maximum oil content, oil yield and crude protein content in both the crops. This was followed by T<sub>7</sub> (75 % S ha<sup>-1</sup> through gypsum + 25 % S ha<sup>-1</sup> through poultry manure) in both the crops. The lowest values for yield, quality, nutrient uptake and nutrient availability were recorded in T<sub>1</sub>-No sulphur / RDF alone (control) in both the crops. T<sub>1</sub> recorded Maximum N,P,K and S availability in both the crops.

**Keywords:** integrated sulphur, nutrient availability, sunflower

### Introduction

The oilseeds form essential part of human diet, besides it produces basic raw materials for agro-based industries. Sunflower has large acreage under various oilseeds in different agro-climatic zones of this country. The average Indian consumer uses relatively lesser quantities of edible oil, no doubt influenced by this modest level of income. Sunflower (*Helianthus annuus* L.) holds great promise as an oilseed crop because of its short duration, photo- in – sensitivity and wide adaptability to different agro-climatic regions and soil types. Sunflower seed contains about 48-53 percent edible oil, sunflower oil is a rich source of linoleic acid (64%) which is good for heart patients. The oil is also used for manufacturing hydrogenated oil. It can be grown at any time of year and can serve as an ideal catch crop during period when the land is otherwise fallow. The existing yield is very low, mainly because of the suboptimal soil fertility. After N,P and K, S is the fourth nutrient, whose deficiency is widespread in India (Sakal *et al.*, 2001) [13]. Sulphur application has many advantages for sunflower regarding growth parameters and yield and quality. Each unit of fertilizer sulphur generates 3-5 units of edible oil, a commodity needed by every family. Sulphur can be rightly called as fourth major element of the plant because it is a constituent of three amino acids viz., Cysteine, Cystine and Methionine which are the building blocks of protein and helps in the formation of chlorophyll and synthesis of oils. Sulphur improves protein and oil percentage in seeds. Sing (1999) [12] reported that application of sulphur increased the uptake of various macro and micro nutrients in groundnut.

Sulphur is required to attain high yield, biological yield, harvesting index and oil content, as affected by biofertilizers and sulphur application. Application of 2 kg S ha<sup>-1</sup> increased seed yield by 38%.

Ever since the role of soil fertility in crop production has been recognized, the use of organic manures has become an imperative need for successful farming. It is a known fact that organic matter provides considerable amount of both macro and micronutrients. It improves soil structure through increased aggregation, which favourably influences the tilth, water infiltration, moisture retention, drainage, aeration, temperature and root penetration besides prevention of crusting. Utilization of all possible sources of organic matter such as vermicompost, FYM, poultry manure helps in improving the soil fertility status and also enhances the yield of oilseed crops. In addition, integration of organics with inorganic improves the physiological system of crop growth, provides adequate growth regulating substances, modifies physico-chemical properties of soil and thus augmenting crop yields. With this background the investigation was carried to increase sunflower yield and quality through integrated sulphur management.

### Material and Methods

Field investigations were carried out during March 2012 and August 2012 at Experimental Farm, Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalai Nagar. The soil of experimental field was clay loam in texture. The soil was low in available Nitrogen, medium in available Phosphorous, high in available Potassium and low in available Sulphur. The sunflower cv. Sumbred was chosen for the study. The experiment was laid out in Randomized block design with three replications. The experiment consisted of eleven treatments viz., T<sub>1</sub> – No sulphur / RDF alone (control), T<sub>2</sub> – 45 kg S ha<sup>-1</sup> through gypsum, T<sub>3</sub> – 45 kg S ha<sup>-1</sup> through vermicompost, T<sub>4</sub> – 45 kg S ha<sup>-1</sup> through poultry manure, T<sub>5</sub> – 45 kg S ha<sup>-1</sup> through lignite flyash, T<sub>6</sub> – 75 % S ha<sup>-1</sup> through gypsum + 25 % S ha<sup>-1</sup> through vermicompost, T<sub>7</sub> – 75 % S ha<sup>-1</sup> through gypsum + 25 % S ha<sup>-1</sup> through poultry manure, T<sub>8</sub> – 75 % S ha<sup>-1</sup> through gypsum + 25 % S ha<sup>-1</sup> through lignite flyash, T<sub>9</sub> – 50 % S ha<sup>-1</sup> through gypsum + 50 % S ha<sup>-1</sup> through vermicompost, T<sub>10</sub> – 50% S ha<sup>-1</sup> through gypsum + 50% S ha<sup>-1</sup> through poultry manure and T<sub>11</sub>– 50% S ha<sup>-1</sup> through gypsum + 50 % S ha<sup>-1</sup> through lignite flyash. The recommended dose of 60:90:60 kgs of NPK ha<sup>-1</sup> was applied in the form of urea, DAP and muriate of potash. Sulphur @ 20 kg ha<sup>-1</sup> was applied through Gypsum as per the treatments.

### Results and Discussion

#### Yield attributes (Table 1)

The integrated sulphur management practices significantly influenced the yield attributes viz., head diameter, total number of seeds head<sup>-1</sup>, number of filled seeds head<sup>-1</sup>, seed filling percent and 100 seed weight in both the crops.

Application of 75% S through gypsum + 25 % S ha<sup>-1</sup> through vermicompost (T<sub>6</sub>) significantly increased the head diameter, total number of seeds head<sup>-1</sup>, number of filled seeds head<sup>-1</sup>, seed filling percent and 100 seed weight and seed and stalk yield over the other treatments. Sulphur is known to play a vital role in the formation of aminoacids. It had favourable effect on yield attributes due to proper partitioning of photosynthates from source to sink. These findings were earlier reported by Syed Shajat Hussain *et al.* (2011) [14]. The

least values for yield attributes were recorded under T<sub>1</sub> (No sulphur – RDF alone), could be due to poor availability of S and other nutrients. These findings were earlier reported by Poonkodi and Poomurugesan (2005) [8]. The increase in yield under this treatment might be due to significant increase in yield attributes leads to seed and stalk yield. This might be due to the influential role played by sulphur in increasing both growth yield attributes. Supply of S in addition to N,P and K might be the lifting factor behind the increased seed and stalk yield (Kapilashkehawat and Shivay, 2008) [6]. T<sub>1</sub> (Control) recorded lesser seed and stalk yield in both the crops. This might be due to absence of sulphur resulted in reduced growth and yield attributing characters timely seed and stalk yield. These findings are in line with Ravikumar (2001) [9].

#### Yield (Table 1)

All the integrated sulphur management practices significantly influenced the seed and stalk yield of sunflower in both the crops over sulphur without integration of organics. Among the treatments tried, T<sub>6</sub> (75% S through gypsum + 25 % S ha<sup>-1</sup> through vermicompost) significantly recorded higher seed yield of 1825.67 and 1927.39 kg ha<sup>-1</sup> and stalk yield of 3800.63 and 3898.59 kg ha<sup>-1</sup> in first and second crop respectively. This treatment registered higher seed yield over no sulphur application (T<sub>1</sub>) and the yield increase being 44.18 and 48.01 per cent respectively in first and second crop respectively. Sulphur application increased the chlorophyll content in leaf and gave a significant positive correlation between chlorophyll content in leaf and crop yield (Sinha *et al.*, 1995) [13]. Control treatment T<sub>1</sub> (No sulphur – RDF alone) recorded lesser seed and stalk yield in both the crops. These findings are in line with Ravikumar *et al.* (2001) [9].

#### Quality characters (Table 2)

The integrated sulphur management practices significantly increased the oil content, oil yield and crude protein content. The maximum oil content of 38.42 and 38.65 percent was recorded with 75% S through gypsum + 25 % S ha<sup>-1</sup> through vermicompost (T<sub>6</sub>) and least value was recorded under no S applied plot (T<sub>1</sub>). Increase in oil content by sulphur application might be attributed to involvement of sulphur in the biosynthesis of oil (Mudd, 1967) [7]. Lowest values of oil and protein content in sunflower seeds was observed in the treatment T<sub>1</sub> (No sulphur) in both the crops. This might be due to lesser availability and uptake of nutrients for the oil and protein synthesis in the crop. Similar view was expressed by Renugadevi and Balamurugan (2002) [10].

#### Nutrient uptake (Table 2)

In both the crops, N,P,K and S uptake were significantly influenced by integrated sulphur management practices. Among the treatments tried, T<sub>6</sub> registered the higher amount of N,P,K and S uptake in both the crops respectively. This might be due to optimum rate of 'S' application through gypsum along with vermicompost increased the uptake of N,P,K & S and ultimately more utilization of these nutrients, which in turn enhanced their concentration and uptake (Bhagat *et al.*, 2003) [1]. Increase in the uptake of N,P and K, S by sunflower with gypsum application was due to the combined effect of increase in yield and nutrient content in plants. Similar results were earlier reported by Devakumar and Gajendra Giri (1990) [2]. T<sub>1</sub> recorded minimum values for N,P, K & S uptake due to lesser availability of nutrients in both the crops.

**Table 1:** Effect of integrated sulphur management on yield attributes and yield of sunflower

Treatments	Head Diameter		Total No. of Seeds head <sup>-1</sup>		No. of filled seeds head <sup>-1</sup>		Seed filling %		100 seed at (g)		Seed yield (Kg ha <sup>-1</sup> )		Stalk Yield (kg ha <sup>-1</sup> )	
	I Crop	II Crop	I Crop	II Crop	I Crop	II Crop	I Crop	II Crop	I Crop	II Crop	I Crop	II Crop	I Crop	II Crop
T <sub>1</sub>	12.60	13.42	489.47	534.35	310.35	337.28	63.40	63.86	3.83	3.90	1266	1302	2872	2957
T <sub>2</sub>	14.50	15.32	608.39	628.31	442.17	460.11	72.67	73.22	4.85	4.92	1514	1560	3489	3564
T <sub>3</sub>	14.27	15.11	579.27	585.25	405.15	421.13	69.94	71.95	4.68	4.68	1467	1511	3431	3499
T <sub>4</sub>	13.98	14.81	545.14	550.16	371.12	390.07	68.07	70.90	4.45	4.43	1399	1435	3385	3435
T <sub>5</sub>	13.50	14.45	515.12	523.08	342.07	361.03	66.40	69.02	4.20	4.21	1371	1403	3327	3371
T <sub>6</sub>	18.67	19.72	860.58	934.37	714.68	787.35	83.04	84.26	6.20	6.42	1825	1927	3800	3898
T <sub>7</sub>	17.45	18.53	778.06	815.26	634.45	672.28	81.54	82.15	5.74	5.87	1765	1850	3735	3805
T <sub>8</sub>	15.50	16.58	677.38	705.23	520.19	548.12	76.79	77.72	5.33	5.36	1656	1721	3602	3683
T <sub>9</sub>	16.37	17.44	725.16	757.21	568.27	601.21	78.36	79.39	5.52	5.60	1702	1780	3668	3746
T <sub>10</sub>	14.85	15.77	658.47	687.25	500.47	525.29	76.00	76.43	5.28	5.32	1601	1654	3590	3662
T <sub>11</sub>	14.68	15.56	642.99	673.19	486.25	512.18	75.62	76.08	5.16	5.14	1555	1613	3540	3595
S.Ed	0.30	0.36	7.75	7.92	7.59	8.42	0.20	0.22	0.08	0.09	24	35	27	31
CD (P = 0.05)	0.61	0.73	15.50	15.84	15.18	16.85	0.41	0.45	0.16	0.19	48	25	55	62

**Table 2:** Effect of integrated sulphur management of Quality parameters and Nutrient uptake of sunflower

Treatments	Oil Content (%)		Crude Protein Content (%)		Nutrient uptake (kg ha <sup>-1</sup> )							
	I Crop	II Crop	I Crop	II Crop	N		P		K		S	
					I Crop	II Crop	I Crop	II Crop	I Crop	II Crop	I Crop	II Crop
T <sub>1</sub>	37.15	37.21	20.48	21.37	67.52	68.43	14.33	15.42	61.48	62.27	6.71	7.38
T <sub>2</sub>	37.63	37.69	22.58	23.44	71.34	72.23	15.89	16.79	64.77	65.46	7.93	8.61
T <sub>3</sub>	37.50	37.55	22.25	22.86	70.88	71.66	15.67	16.49	64.29	65.25	7.63	8.36
T <sub>4</sub>	37.38	37.41	21.39	22.47	70.29	71.36	15.39	16.21	63.86	64.75	7.33	8.16
T <sub>5</sub>	37.22	37.27	21.67	21.99	69.86	70.92	15.12	15.94	63.37	64.22	7.12	7.95
T <sub>6</sub>	38.42	38.65	27.53	28.39	77.62	78.32	19.42	20.21	70.45	71.27	11.47	12.25
T <sub>7</sub>	38.27	38.32	26.21	27.29	76.35	77.23	18.37	19.20	69.31	70.19	10.38	11.21
T <sub>8</sub>	38.07	38.21	24.19	25.22	74.16	75.20	16.50	17.30	67.18	68.17	8.73	9.28
T <sub>9</sub>	38.16	38.15	25.18	26.23	75.25	76.13	17.42	18.25	68.29	69.18	9.54	10.18
T <sub>10</sub>	37.35	37.78	23.68	24.55	72.68	73.36	16.41	17.22	65.78	66.47	8.58	9.10
T <sub>11</sub>	37.76	37.67	23.52	24.41	71.87	72.63	16.14	17.05	65.26	66.16	8.22	8.87
SEd	0.05	0.06	0.07	0.08	0.43	0.43	0.17	0.18	0.38	0.39	0.20	0.21
CD (P = 0.05)	0.10	0.13	0.15	0.17	0.86	0.86	0.35	0.36	0.76	0.78	0.41	0.42

**Table 3:** Effect on integrated sulphur management of nutrient uptake on sunflower

Treatments	Nutrient availability (Kg ha <sup>-1</sup> )							
	N		P		K		S	
	I Crop	II Crop	I Crop	II Crop	I Crop	II Crop	I Crop	II Crop
T <sub>1</sub>	237.20	241.22	22.45	27.61	343.64	349.70	22.60	29.15
T <sub>2</sub>	224.26	227.29	20.49	24.31	338.31	335.41	19.40	25.30
T <sub>3</sub>	226.15	230.42	20.21	25.21	339.25	338.22	20.35	26.27
T <sub>4</sub>	229.21	233.31	21.45	26.32	340.42	342.19	21.41	27.10
T <sub>5</sub>	230.13	239.28	21.20	27.15	342.27	345.23	22.27	28.23
T <sub>6</sub>	191.50	201.20	15.28	20.40	305.11	310.23	13.08	17.22
T <sub>7</sub>	197.37	207.31	16.19	21.37	315.21	313.32	14.20	19.31
T <sub>8</sub>	205.29	213.39	17.21	22.30	320.29	319.29	15.31	21.20
T <sub>9</sub>	213.19	219.28	18.32	21.42	329.27	322.30	16.42	22.19
T <sub>10</sub>	222.69	221.19	19.46	22.37	334.35	325.41	17.29	23.31
T <sub>11</sub>	223.48	225.31	20.29	23.25	335.22	330.28	18.37	24.28
SEd	0.53	0.49	0.55	0.30	0.53	0.48	0.78	0.78
CD (P = 0.05)	1.12	1.03	1.15	0.64	1.10	1.01	1.64	1.64

**Post harvest soil available nutrient status (Table 3)**

The integrated sulphur management practices influenced the post harvest nutrient status *viz.*, N, P, K & S over the treatments received sulphur without organics. Among the various practices tried, T<sub>1</sub> recorded higher values for soil available N,P, K & S in both the crops. This might be due to poor uptake of nutrients by the crop as result of lesser foraging capacity. Similar results were also reported by Kalaiyarasan (2000) [5]. N,P, K & S were low in T<sub>6</sub> due to more uptake of nutrients by the crop (Gandhi, 2011) [3].

**References**

- Bhagat GJ, Abdul Hamid SP, Bonda MD Giri, Sajid M. Effect of irrigation and sulphur on growth and yield attributes of rabi sunflower. Res. On crops. 2003; 4(1):56-59.
- Devakumar M, Gajendra Giri. Effect of weed control and gypsum application on uptake of N,P, Ca and S by groundnut and weeds. Ind. J Agron. 1990; 44(2):400-403.
- Gandhi G. Study on response of sunflower genotypes to sulphur fertilization grown under veeranamayacut regions. M.Sc. (Ag.). Thesis, Annamalai University, Annamalai Nagar, 2011.

4. Jeyabal A. Agronomic evaluation of bio-digested slurry in rice-blackgram cropping system. Ph.D., Thesis, Annamalai University, Annamalai Nagar, 1996.
5. Kalaiyarasan C. Studies on sulphur management in groundnut. CV. VRI 2. M.Sc. (Ag.) Thesis, Annamalai University, Annamalai Nagar, 2000.
6. Kapilashkehawat, Shivay YS. Effect of Boron- sulphur interaction on their uptake and quality parameters of mustard (*Brassica juncea* L.) and sunflower (*Helianthus annus* L.). J Indian Soci. Sci. 2008; 56(2):225-230.
7. Mudd JB. Fat Metabolism in plants, Ann. Plant Physiol. 1967; 18:229-252
8. Poonkodi P, Poomurugesan AV. Effect of sulphur on growth and yield of sunflower (*Helianthus annus* L.). Andhra Agric. J. 2005; 52(3&4):600-601.
9. Ravikumar C. Studies on sulphur management in sunflower. M.Sc. (Ag.) Thesis, Annamalai University, Annamalai Nagar, 2001.
10. Renugadevi J, Balamurugan P. Effect of biofertilizer in seed yield and seed quality in sunflower. Madras Agric. J. 2002; 89(1-3):135-137.
11. Sarkar RK, Mallic RB. Effect of Nitrogen, sulphur and foliar spray of nitrate salts on performance of spring sunflower (*Helianthus annus* L.). Indian J Agric Sci. 2009; 79(12):986-90.
12. Sing RA. Nitrogen, sulphur and calcium relationship in sustainable production of Indian mustard (*Brassica juncea*) on denuded land. Indian J Agron. 1999; 44(4):820-825.
13. Sinha RB, Sakal P, Kumar S. Sulphur and phosphorus nutrition of winter maize in calcareous soil. J Ind. Soc. Soil Sci. 1995; 43(3):413-418.
14. Syed Shajat Hussain, F.A. Misger, Amit Kumar and Bala MH. Response of nitrogen and sulphur on biological and economic yield of sunflower (*Helianthus annus* L.). Research J Agric. Sci. 1995, 2011; 2(22):308-310.