



P-ISSN: 2349-8528  
 E-ISSN: 2321-4902  
 IJCS 2019; 7(5): 3302-3305  
 © 2019 IJCS  
 Received: 18-07-2019  
 Accepted: 21-08-2019

**Shreya Singh**  
 M.Sc. Student, School for Home  
 Science F.S.T B.B.A.U,  
 Lucknow, Uttar Pradesh, India

**Neetu Singh**  
 Assistant Professor, School for  
 Home Science F.S.T B.B.A.U,  
 Lucknow School for Home  
 Science, Uttar Pradesh, India

**Corresponding Author:**  
**Shreya Singh**  
 M.Sc. Student, School for Home  
 Science F.S.T B.B.A.U,  
 Lucknow, Uttar Pradesh, India

## International Journal of Chemical Studies

### Development of probiotics & prebiotics product as weaning food for infants

**Shreya Singh and Neetu Singh**

#### Abstract

The present investigation entitled “Development of probiotics & prebiotics product as weaning food for infants” was based upon Probiotics and prebiotics which is beneficial for bacteria grow back and good digestion capability. Prebiotics function is to increase our gut healthy bacteria groups such as bifid bacterium and lactobacillus. Food containing probiotics (bacteria) and prebiotics (fermented fiber) are called symbiotic food because a component stimulates the activity of the other, being twice effective. In this research work mainly three weaning food were developed with different compositional values of milk and starch weaning food i.e. Milk (120ml), starch (30ml), Milk and bread weaning food, Milk (120ml), bread (3 slice), Milk, fruits pulp weaning food, milk (120ml), fruits pulp (56g). Highly accepted weaning food is milk and starch weaning food among the three products.

**Keywords:** Weaning food, digestion, stimulates, probiotics and capability

#### Introduction

Probiotics and prebiotics can help the beneficial bacteria grow back. A person needs a balance of intestinal bacteria favorable to digestion in order for your body, especially the intestines, to utilize minerals and other nutrients from your food. This bacteria works off carbohydrates and sugars, often called fructooligosaccharides, found in vegetables and fruits, a large part of a balance diet. A balance diet is essential for health to provide the nutrients needed for energy and vitality and it consists of protein, carbohydrates, fiber, vitamins and minerals, and fats. It may come as a surprise, but not all bacteria are bad for us. It's common to hear new report about harmful bacteria (pathogens) these days, & most of us are familiar with some of their names-*Listeria monocytes*, *clostridium botulinum* and various strains of salmonella, for instance. But other families of bacteria are not injurious to health and may in fact; provide some health benefits, particularly those in the *lactobacillus* and *Bifidobacterium* genera. Probiotics are “live microorganism which when administered in adequate amounts confer a beneficial health effect on the host.” These micro-organisms do not promote or cause disease. Gastrointestinal flora influence health, but the composition of flora can be adapted by consuming prebiotics or probiotics. Prebiotics are no digestible substances that stimulate the growth of health-promoting bacteria in the colon, such as bifid bacteria. They are live microbial feed supplements which improve the intestinal microbial balance. Immediately after birth, bacterial colonization of the infant's gut begins with bifid bacteria and lactobacilli. These organisms are transferred from the maternal microbial flora in the genital tract and colon during delivery, and also from the environment. Gut-associated immune tissue comprises 80% of the immune system, making the composition of intestinal flora an important factor in the immune system. It is believed that the onset of many diseases possibly relates to disruption of the early colonization of the gut. many Preterm infants have delayed colonization of the gut with beneficial flora by three to four weeks, but colonization with pathogens occurs earlier and contributes to health problems, such as necrotizing enter colitis (NEC).

#### Materials and Methods

The present investigation entitled “Development of probiotics & prebiotics product as weaning food for infants” was carried out to outlay the effect of probiotics as well as prebiotics among infants under the following heads:

1. Prebiotic and probiotic property of the sample
2. Weaning food and its standardization

### Selection of ingredient for weaning food

Weaning food is a food for infant (6 months to 2 year), it is a source of food other than mother's breast milk. Weaning is the introduction of solid food in a baby's diet during the first year after birth. Starchy foods (rice, bread, fruits) are rich in prebiotic to enhance or activate the growth of probiotics are mixed in milk, curd is used as a probiotic for the fermentation. Probiotic are the microorganism that are present in curd and live in gut of humans and help in the digestion.

### Nutritional property of taken sample

This step involved nutritional analysis of the milk involving different parameters like-

Determination of fat percentage, estimation of protein percentage, amount of carbohydrate and level of vitamin C percentage.

**Tools:** For the characterization of nutritional profile of milk, curd, rice, fruits and bread DIET. CAL. Software is used.

**Product development:** This phase involved the all method of production of weaning food by using different ingredients in milk like-milk with bread, milk with starch and milk with fruits pulp.

### Results and Discussion

#### Prebiotic and Probiotic Property of the Sample

Prebiotics Function is to increase our gut healthy bacteria groups such as bifidobacterium and lactobacillus. Food containing probiotics (bacteria) and prebiotics (fermented fiber) are called symbiotic food because a component stimulates the activity of the other, being twice effective. This fermentation process feeds beneficial bacteria colonies (including probiotic bacteria) and helps to increase the number of desirable bacteria in our digestive systems that are associated with better health and reduce disease risk.

The colonic micro flora is important to health. The growth and metabolism of the many individual bacterial species inhabiting the large bowel depend primarily on the substrates available to them, most of which come from the diet [1]. This has led to attempts to modify the structure and metabolic activities of the community through diet-using probiotics and prebiotics. Probiotics are live microbial food supplements. The best known are the lactic acid bacteria and bifidobacteria, which are widely used in yoghurts and other dairy products. These organisms are non-pathogenic and non-toxic, retain viability during storage, and survive passage through the stomach and small bowel. Prebiotics are non-digestible food ingredients which selectively stimulate the growth or activities, or both, of lactobacilli or bifid bacteria in the colon, thereby improving health.

#### Nutritional value of milk

The table given below is the interpretation of nutritional analysis of milk value give in 100g.

**Table 1:** Product and Carbohydrate energy calcium

Product name	Protein (g)	Fat (g)	Carbohydrate (g)	Energy (k. Call)	Calcium (mg)
Milk buffalo's (100 ml)	4.3	6.5	5.0	117	210
Milk buffalo's (120ml)	5.2	7.8	6	140	252
Milk cow's	3.2	4.1	4.4	67	120

#### Nutritional characteristics of rice value give in 100 g-

The table given below is the interpretation of nutritional

analysis of rice.

**Table 2:** Protien energy fat Carbohydrate calcium

Product	Protein (g)	Energy (k. call)	Fat (g)	Carbohydrate (g)	Calcium (mg)
Raw rice (100g)	6.8	370	0.5	81.7	11.0
Raw rice (30g)	1.3	103.5	0.2	23.46	2.6
Rice milled	6.8	345	0.5	83	10

### Develop weaning food

#### Preparation of weaning food with milk and starch

Quantity used for the milk and starch weaning food were- Required amount of milk=120ml required amount of rice starch=30ml required amount of curd =1/2 tbsp. It enhances the digestibility of infant because it provides prebiotics and probiotics.

#### Prepared weaning food with milk and fruits pulp

Quantity used for the milk and fruits pulp weaning food were- Required amount of milk = 120ml required amount of fruits pulp (banana & muskmelon) = 56g required amount of curd for fermentation=1/2 tbsp.

#### Prepared weaning food with milk and bread

Quantity used for the milk and bread weaning food were required amount of milk=120ml, required amount of bread =3 slice and required amount of curd for fermentation =1/2 tbsp.

### Summary and Conclusion

#### Summary

Prebiotics function is to increase our gut healthy bacteria groups such as bifidobacterium and lactobacillus. Food containing probiotics (bacteria) and prebiotics (fermented fiber) are called symbiotic food because a component stimulates the activity of the other, being twice effective. This fermentation process feeds beneficial bacteria colonies (including probiotic bacteria) and helps to increase the number of desirable bacteria in our digestive systems that are associated with better health, reduce disease and better digestive system.

In the preparation of milk + starch weaning food, the quantity of milk taken was (120ml) and the starch quantity (30ml) In the preparation of milk + bread weaning food, the quantity of milk taken was (120ml) and the bread quantity (3 slices). Preparation of milk + fruits pulp weaning food, the quantity of milk taken was (120 ml) and the fruit pulp quantity (56g).

#### Conclusion

In this research work mainly three weaning food were developed with different compositional values of milk and starch weaning food i.e. Milk (120ml), starch (30ml), Milk and bread weaning food, Milk (120ml), bread (3 slice), Milk, fruits pulp weaning food, milk (120ml), fruits pulp (56g). Highly accepted weaning food is milk and starch weaning food among the three products. Weaning is mainly recommended to infant (6 months to 2 year) their protein, carbohydrate, fat and vitamin C. Effective in gums and different digestive related problems.

#### References

1. Aalander M, Matto J, Kneifel W, Johansson M, Kogler B, Crittenden R. Effect of galacto-oligosaccharide supplementation on human fecal microflora and on survival and persistence of Bifidobacterium Lactis Bb-12 in the gastrointestinal tract. International Dairy Journal.

- 2001; 11:817-825.
2. Adam A, Levrat-Verny MA, Lopez HW, Leuillet M, Demigne C, Remesy C. Whole wheat and triticale flours with differing viscosities stimulate cercal fermentations and lower plasma and hepatic lipid in rats. *Journal of Nutrition*. 2001; 131:1770-1776.
  3. Anderson DM. From Pediatric Nutrition. *Handbook of Pediatric Nutrition*. Edited by: Samour PQ, Helm KK. James and Bartlett Publishers, Sudbury, Massachusetts, 2005, 53-71.
  4. Gibson GR, Beatty EH, Wang X, Cummings JH. Selective stimulation of bifidobacteria in the human colon by oligofructose and inulin. *Gastroenterology*. 1995; 106:975-82.
  5. Nutritional advantages of probiotics and prebiotics. Marteau P, Boutron-Ruault MC. *British Journal of Nutrition*. 2002; 87(2):5153-5157. DOI: 10.1079/BJN2002531.
  6. Probiotic bacteria in fermented foods: product characteristics and starter organisms, Knut J Heller *Probiotics and Prebiotics-a report by Marry Ellen Sanders, PhD and Glenn Gibson, Ph.D. Consultant, Dairy and Food Culture Technologies, and Professor of Food Microbiology, University of reading.*
  7. Probiotics and prebiotics: can regulating the activities of intestinal bacteria benefit health George T Macfarlane, John H Cummings Probiotics, prebiotics and synbiotics- a review, Kavita R Pandey, Suresh R Naik, Babu V Vakil.
  8. Robles Alonso V, Guarner F. Linking the gut microbiota to human health. *British Journal of Nutrition*. 2013; 109:S21-S26.
  9. Sanders ME *et al.* An update on the use and investigation of probiotics in health and disease. *Gut*. 2013; 62(5):787-796.
  10. Wu GD, Lewis JD. Analysis of the human gut microbiome and association with disease. *Clinical Gastroenterology Hepatology*. 2013; 11(7):774-777.