



P-ISSN: 2349-8528

E-ISSN: 2321-4902

IJCS 2019; 7(6): 2278-2282

© 2019 IJCS

Received: 01-09-2019

Accepted: 03-10-2019

AM Shinde

Department of Soil Science and
Agricultural Chemistry, Post
Graduate Institute, Dr. PDKV,
Akola, Maharashtra, India

MD Sarode

Department of Soil Science and
Agricultural Chemistry, Post
Graduate Institute, Dr. PDKV,
Akola, Maharashtra, India

AR Deshmukh

Department of Soil Science and
Agricultural Chemistry, Post
Graduate Institute, Dr. PDKV,
Akola, Maharashtra, India

Assessment of fertility status of Babhulgao block of the central research station Dr. PDKV Akola

AM Shinde, MD Sarode and AR Deshmukh

Abstract

Soil of Babhulgao block of Central research station Dr. PDKV Akola was investigated for the chemical properties and Macronutrient or micronutrient status of surface soils. The exact position of soils sampling site was determined with the help GPS.

All the soils under study were slightly alkaline to moderately alkaline in reaction, EC value for these soils within safe limit. The organic carbon content in these soils is low to medium and soil of farms is moderately calcareous to calcareous due to presence of CaCO_3 .

The soils of Babhulgao block contain 146.76 to 213.25 kg ha^{-1} available nitrogen, 14.52 to 18.82 kg ha^{-1} available phosphorus and 247.8 to 341.6 kg ha^{-1} available potassium. The soils of babhulgao block contain available micronutrients i.e. Zn, Cu, Fe, Mn in the range of contains 0.33 to 0.69 mg kg^{-1} , 0.28 to 3.28 mg kg^{-1} , 2.32 to 7.80 mg kg^{-1} , 1.50 to 8.60 mg kg^{-1} respectively, Zn, Cu, Fe, Mn respectively.

The nutrient index value calculated for soils of Babhulgao block shows that soils has moderate NIV for organic carbon, low NIV for available nitrogen and moderate for phosphorus, high NIV for available potassium. The micronutrients status show high NIV for iron, copper and manganese, The Babhulgao block shows the low NIV for zinc.

It is revealed that in the soils of Babhulgao block, the availability of nitrogen, iron, zinc decreases with increase in CaCO_3 content. Available copper also decreases with increase in soil pH. The availability of zinc decrease due to increase in EC. The availability of manganese, copper, increase with increase in organic carbon status in course of investigation.

Keywords: Babhulgaon, fertility

1. Introduction

The area of soil science research which is most directly related to agricultural productivity, soil fertility and fertilizer use research and this is also an area where the expectations are high. All researches in soil fertility have one common goal that is to assess nutrient supplying capacity of the soil, deficiencies of nutrient if any and to supply nutrient based on crop needs. Thus, in the game of crop production, there are three dependent and yet interdependent players the soil, the plant and fertilizers, each one of them key players (Goswami, 1999) ^[1]. India's population is variously projected at 1330 million to 1620 million by 2020 (16-17 million population added each year) and food grain demand by 2020 is estimated at 260-300 million tons (117 million tons of rice, 93 million tons of wheat, 28 million tones of course grain and 24 million tones of pulses). The challenge during the next millennium is to achieve and sustain growth rates high enough to feed the swelling population without degrading the environment (NAAS, 1997).

Indian soils have been developed under different climatic conditions such as semiarid, tropical and sub-tropical and thus vastly differ in their properties. Research work done so far is still inadequate to decide their fertility status. In India, the black soils are termed as black cotton soils. These soils are characterized by dark greyish brown to very dark color, high montmorillonitic clay content, high coefficient of expansion and shrinkage and show deep cracking during dry seasons. They have varying depth ranging from shallow to very deep. They cover extensive areas in the Deccan plateau as well as Central India. Black and associated soils are developed from Deccan trap and early Pleistocene alluvial deposits (Flood plain) under tropical monsoonic environment occur extensively in Vidarbha region of Maharashtra.

Swell-shrink soil series of Maharashtra state were studied during the year 1991-1992 with respect to their available major and micronutrient status. The result shows that these soils were

Corresponding Author:**AM Shinde**

Department of Soil Science and
Agricultural Chemistry, Post
Graduate Institute, Dr. PDKV,
Akola, Maharashtra, India

low in available Nitrogen, very low to moderate in Phosphorus and moderate to very high in Potassium contents. The DTPA extractable Fe, Mn, Zn and Cu were in the range of 6.1 to 26, 14.6 to 63.0, 0.58 to 1.70 and 1.7 to 6.1 PPM, respectively. Considering the critical limits of these micronutrients except zinc, all other micronutrients were well supplied. The DTPA Fe was negatively correlated with pH, organic carbon, clay, CEC and CaCO₃. The available Mn was negatively but significantly correlated with pH and CaCO₃, while Zn showed positive relationships with organic carbon and clay content of the soil (Patil and Sonar., 1994). Available boron ranged between 0.04 to 0.85 PPM. Available boron shows positive and significant correlation with clay, silt, pH, EC, organic carbon and available N and K where as negative correlation with available P and sand content of soil (Deshmukh *et al.*, 2007) [3].

Regarding the nutrient status, all soils were low in organic carbon and available nitrogen, however, medium in available phosphorous and high to very high in available potassium. The micronutrient delineation of these soils were marginal, adequate, high and low to marginal in Fe, Mn, Cu and Zn respectively. The availability of micronutrient in Fe and available Cu were significantly and positively correlated among themselves. Available Mn and other micronutrient did not show any significant relationship between themselves. (Jibhate *et al.*, 2009) [5]

2. Material and Methods

The experiment was laid out at Babhulgao of the central research station Dr. PDKV, Akola in the year 2015-2016 with an objective to assess the fertility status and to develop the fertility index of soils of the farms and prepare soil fertility map".

Climate, geology, vegetation, topography and time are five factor's which influence the formation of soils. The geology of the district is transitional with dominance of Deccan basalts in Western portion, sand stones and shales in Eastern portion. Different geological formations met in the vicinity of eastern part of the central research station. Effect of climate has been

modified by relief rendering different soil climate and subsequent local arid and humid conditions. Due to undulating topography, soil erosion has also played on down a slope.

Most of the area under cultivation comprises of black soils. Katepurna is the main river flowing through the district.

Two type of soils have been observed in the district namely medium black soil occurring in plain central part of trap origin and deep black soil occurring in valley in northern part, predominantly laying on pediment alluvial plains and flood plains to transitional slopes. Being moderately deep and clayey of montmorillonitic mineralogy they are more water retentive.

The climate of the region is arid monsoonic type and characterized by the three distinct season *viz.* with hot and dry weather from March to May, monsoon warm and rainy from June to October and winter dry mid cold from November to February.

The central research station is situated about 2 km east of Akola town. It is located at longitude 77°02'44" to 77°04', 59"E, latitude 20°42'15" to 20°43'18"N.

Average rainfall is about 1192.5mm. Distribute in 48 rainy days (30 years recorded) in the year. About 77.7 percent of the mean total rainfall is received during the monsoon period in 40 rainy days (June-September) about 3 percent during the post monsoonic period.

The mean annual maximum and minimum temperature are 30 degree Celsius and 24.4 degree Celsius respectively. Summer month are fairly hot with maximum air temperature ranging from 37 to 44 degree Celsius with relative humidity ranging from 26 to 29 percent. April and May are the hottest months of the year (40.5 to 46.4 degree Celsius) with lowest relative humidity in April. The winter months experience mild cold with average temperatures ranging from 21 to 24 degree Celsius.

2.1 Soil nutrient index and preparation of thematic maps

Soil nutrient index was calculated as per six tier system as follows (Ramamurthy and Bajaj, 1969) [6]

$$\text{Nutrient Index Value} = \frac{(\text{No. of samples in very low} \times 0.5) + (\text{No. samples in low} \times 1.0) + (\text{No. samples in medium} \times 1.5) + (\text{No. of samples in moderately high} \times 2) + (\text{No. of samples in high} \times 2.5) + (\text{No. samples in Very high} \times 3)}{\text{No. of Samples taken}}$$

Table 1: Ratings of Nutrient index value for major nutrients

Sr. No	Category	Value
1	Very low	0.5 to 0.75
2	Low	0.76 to 1.25
3	Moderate	1.26 to 1.75
4	Moderately high	1.76 to 2.25
5	High	2.26 to 2.75
6	Very high	2.76 to 3.00

Table 2: Ratings for major nutrients

Sr. No	Nutrient element	Category					
		Very low	Low	Medium	Moderately high	High	Very high
1	Organic Carbon (%)	<0.20	0.21-0.40	0.41-0.60	0.61-0.80	0.81 -1.0	>1.0
2	Available N (kg ha ⁻¹)	<140	141-280	281-420	421-560	561-700	>700
3	Available P (kg ha ⁻¹)	<7.0	7.1-14.0	14.1-21.0	21.1-28.0	28.1-35.0	>35
4	Available K (kg ha ⁻¹)	<100	101-150	151-200	201-250	251-300	>300

Nutrient index for available micronutrient was calculated as per formula given by Ramamurthy and Bajaj (1969) [6]

$$\text{Nutrient Index Value} = \frac{(\text{No. of samples in low} \times 1) + (\text{No. of samples in medium} \times 2) + (\text{No. of Samples in high} \times 3)}{\text{No. of Samples taken}}$$

Table 3: Ratings of Nutrient index value for micronutrients

Sr. No	Category	Value
1	Low	<1.67
2	Medium	1.67-2.33
3	High	>2.33

Table 4: Ratings for available micronutrient

Sr. No	Category		
	Low	Medium	High
DTPA-Fe (mg kg ⁻¹)	<2.5	2.5-4.5	>4.5
DTPA-Mn (mg kg ⁻¹)	<1.0	1.0-2.0	>2.0
DTPA-Zn (mg kg ⁻¹)	<0.6	0.6-1.2	>1.2
DTPA-Cu (mg kg ⁻¹)	<0.2	0.2	>0.2

Table 5: Chemical properties of soils of Babhulgao block

Grid No	PH	EC (dSm-1)	OC (g/kg)	CaCO ₃ (%)	CEC
1	7.9	0.23	4.50	5.81	46
2	7.8	0.27	4.44	5.56	54
3	7.9	0.26	3.00	5.69	52
4	7.8	0.25	3.30	6.44	50
5	7.7	0.20	3.30	5.56	40
6	7.9	0.25	3.00	5.69	50
7	7.8	0.27	4.59	5.31	54
8	7.7	0.24	3.30	6.50	48
9	8.0	0.26	3.30	5.87	52
10	7.7	0.22	3.90	6.44	44
11	7.5	0.27	3.54	6.06	54
12	7.9	0.23	3.00	5.65	46
13	7.7	0.21	3.00	5.83	42
14	8.4	0.23	3.00	6.14	46
15	7.8	0.22	3.90	5.80	44
16	7.8	0.19	4.50	5.67	38
17	8.2	0.20	4.80	5.81	40
18	7.8	0.21	5.10	5.31	42
19	7.7	0.25	3.30	5.00	50
20	7.8	0.31	3.60	5.16	48
21	7.9	0.27	5.70	5.63	54
22	7.9	0.25	5.10	5.14	50
23	7.9	0.22	4.80	5.65	44
24	7.8	0.25	3.30	5.53	50
25	7.8	0.23	5.40	5.25	46
26	7.9	0.21	3.00	5.75	42
27	7.7	0.23	3.90	6.19	46
Mean	7.9	0.25	3.91	5.72	47.11
Range	7.5-8.4	0.19-0.27	3.00-5.70	5.00-6.50	38-54

(For available Fe, Mn, Zn, Cu the critical limits are taken as per given by Katyal and Rattan 2003) [8]. Thematic map of nutrient status of soils of blocks were prepared by using ratings for different nutrients.

2.2 Place/time/duration of research work

The experiment on Babhulgaon of central research station Dr. PDKV, Akola soils initiated during 2015-2016 at central research station Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola. The present investigations have carried out during 2015-2016.

3. Result and Discussion

3.1 Soil nutrient index of available macronutrient for soils of Babhulgaon block.

Table 6: Category wise classification of soil of blocks for available macronutrient of Babhulgaon block soils

Grid no	AV.N	Category	AV.P	Category	AV.K	Category	Oc (g kg-1)	Category
1	200.70	low	16.9	Medium	326	very high	4.5	Medium
2	200.70	low	16.7	Medium	316	very high	4.44	Low
3	150.53	low	15.4	Medium	325	very high	3	Low
4	175.62	low	15.8	Medium	327	very high	3.3	Low
5	175.62	low	18.8	Medium	314	very high	3.3	Low
6	163.07	low	15.9	Medium	302	very high	3	Low
7	163.07	low	17.4	Medium	291	High	4.59	Medium
8	213.25	low	14.6	Medium	336	very high	3.3	Low
9	200.70	low	16.1	Medium	318	very high	3.3	Low
10	175.62	low	17.0	Medium	328	very high	3.9	Low
11	181.89	low	15.8	Medium	342	very high	3.54	Low
12	150.53	low	15.1	Medium	323	very high	3	Low
13	188.16	low	16.8	Medium	288	High	3	Low
14	200.70	low	14.8	Medium	326	very high	3	Low
15	213.25	low	17.0	Medium	288	High	3.9	Low
16	185.65	low	16.8	Medium	318	very high	4.5	Medium
17	213.25	low	15.6	Medium	327	very high	4.8	Medium
18	200.70	low	16.1	Medium	340	very high	5.1	Medium
19	155.55	low	15.2	Medium	317	very high	3.3	Low
20	174.36	low	15.3	Medium	312	very high	3.6	Low
21	200.70	low	17.0	Medium	306	very high	5.7	Medium
22	175.62	low	17.3	Medium	302	very high	5.1	Medium
23	146.76	low	15.1	Medium	289	High	4.8	Low
24	150.53	low	15.2	Medium	292	High	3.3	Low
25	200.70	low	15.6	Medium	291	High	5.4	Medium
26	175.62	low	16.1	Medium	321	very high	3	Low
27	175.62	low	14.5	Medium	330	very high	4.5	Medium
NIV	1	low	2.72	Medium	2.88	High	1.16	moderate

Soils of Babhulgaon block categorized as low to medium for organic carbon content as per six tier system. The soil nutrient index value of Babhulgaon block calculated for organic carbon content on soils is 1.16 g kg⁻¹, Babhulgaon blocks shows the moderate nutrient index for organic carbon content. The nitrogen, phosphorus and potassium are the major essential nutrients and its availability plays very vital role in plant growth and its productivity. The availability of N, P and K govern the fertility status of the soil. The soils of Babhulgaon block contain 146.76 to 213.25 kg ha⁻¹ available nitrogen, 14.52 to 18.8 kg ha⁻¹ available phosphorus and 288 to 340 kg

ha⁻¹ available potassium. As per the six tier classification system of soils for its nutrient content, the soils of Babhulgaon block classified as low for available nitrogen, available phosphorus are classified as medium and high to very high for available potassium. The nutrient index calculated for Babhulgaon block for Available Nitrogen, available phosphorus and available potassium and organic carbon is 1, 2.72, 2.88 and 1.16 respectively

3.2 Soil nutrient index of available micronutrient for soils of Babhulgaon block.

Table 7: Category wise classification of soil of blocks for available micronutrients of Babhulgaon block

Grid No.	Zn (gm kg-1)	Ratings	Cu (gm kg-1)	Ratings	Fe (gm kg-1)	Ratings	Mn (gm kg-1)	Ratings
1	0.58	Low	1.04	High	4.52	High	5.80	High
2	0.40	Low	1.20	High	5.20	High	6.20	High
3	0.51	Low	0.92	High	4.24	High	6.00	High
4	0.52	Low	0.80	High	3.20	High	4.40	High
5	0.33	Low	2.24	High	4.56	High	2.90	High
6	0.60	Low	1.12	High	7.08	High	4.54	High
7	0.55	Low	2.20	High	4.68	High	3.90	High
8	0.44	Low	1.56	High	6.20	High	5.32	High
9	0.64	Medium	1.68	High	4.20	High	1.70	High
10	0.42	Low	0.28	High	2.32	High	4.70	High
11	0.33	Low	1.90	High	4.60	High	3.70	High
12	0.64	Medium	2.64	High	2.48	High	5.91	High
13	0.59	Low	0.68	High	4.56	High	5.41	High
14	0.54	Low	2.40	High	7.36	High	6.39	High
15	0.68	Medium	0.68	High	5.64	High	1.50	High
16	0.59	Low	2.60	High	4.76	High	7.49	High
17	0.53	Low	0.80	High	5.04	High	4.13	High
18	0.49	Low	3.28	High	7.80	High	2.57	High
19	0.55	Low	2.68	High	4.58	High	3.47	High

20	0.66	Medium	1.00	High	4.59	High	3.89	High
21	0.60	Low	2.64	High	5.72	High	4.52	High
22	0.69	Medium	2.64	High	3.80	High	5.60	High
23	0.51	Low	1.20	High	6.40	High	2.00	High
24	0.52	Low	2.84	High	6.68	High	2.90	High
25	0.63	Medium	2.76	High	4.68	High	3.80	High
26	0.48	Low	2.28	High	3.60	High	8.60	High
27	0.60	Medium	0.52	High	4.80	High	5.20	High
NIV	1.25	low	3	High	2.66	High	3	High

The soils of Babhulgao block contain available micronutrients i.e. Zn, Cu, Fe, Mn the range of 0.22-0.69 mg kg⁻¹, 0.28-3.38 mg kg⁻¹, 2.32 to 7.80 mg kg⁻¹, 1.50 to 8.60 mg kg⁻¹ respectively. As per critical limits for available micronutrient given by Katyal J.C. and R.K. Rattan (2003)^[8] the soils are classified for its micronutrient content, the soils of Babhulgao block classified as low to medium for Zinc, medium to high

for Iron and high for manganese and high for available copper. Nutrient index calculated for micronutrients shows the values 1.25 for Zinc, 2.66 for Iron and 3 for both copper and manganese in Babhulgao block. Accordingly the soils of Babhulgao block show low nutrient index value for available zinc, high nutrient index values for available Iron, Manganese and Copper K.

Table 8: Relationship between soil chemical properties with macro and micronutrient for Babhulgao block

1	pH	EC	OC	CaCo3	N	P	K	Zn	Cu	Fe	Mn	S
Ph	1	-	-	-	-	-	-	-	-	-	-	-
EC	-	1	-	-	-	-	-	-	-	-	-	-
OC	-	-	1	-	-	-	-	-	-	-	-	-
CaCo3	-	-	-	1	-	-	-	-	-	-	-	-
N	-	-	0.402*	-	1	-	-	-	-	-	-	-
P	-	-	-	-	-	1	-	-	-	-	-	-
K	-	-	-	0.453*	0.427*	-	1	-	-	-	-	-
Zn	-	-	-	-	-	-446*	-	1	-	-	-	-
Cu	-	-	0.535**	-	-	-	-	-	1	-	-	-
Fe	-	-	-	-	-	-	-	-	-	1	-	-
Mn	-	-	-	-	-	-	-	-	-	-	1	-
S	-	-	-	-	-	-	-	-	-	-	0.455*	1

*Significant at 5% level of significance, **Significant at 1% levels of significance

The significant positive correlation was observed in OC($r=0.402^*$) with available N and Cu ($r=0.535^{**}$). And significant positive correlation between CaCo₃ and available K and significant correlation between available K and available N ($r=0.427^*$) and negative correlation was observed between available P and Zn ($r= -446^*$) and positive correlation observed between Mn and available S ($r= 0.455^*$).

4. Conclusion

1. Soil of Babhulgao, Shivani moderately alkaline in reaction, having EC within the safe limit for crop cultivation. The organic carbon content in these soils is low to medium and CaCO₃ moderately calcareous to calcareous in nature.
2. The soil Babhulgao block low in available nitrogen and medium for available phosphorous and High to Very high in available potassium status.
3. The soils of Babhulgao, blocks medium to high in available Fe, low to medium in available Zn and high in Cu and Mn status.
4. The soil Babhulgao, block, the availability nitrogen, iron, zinc decreases with increase in CaCO₃ content. Available copper also decreases with increase in soil pH. The availability of zinc decrease due to increase in EC.
5. The availability of manganese, copper, increase with increase in organic carbon status.
6. Nutrient index value derived for soils of Babhulgao, block shows that soil has moderate NIV for organic carbon, low NIV for available nitrogen and medium for phosphorous high for available potassium. Babhulgao

block showed low NIV for zinc, and high for iron, manganese and copper.

5. References

1. Goswami NN. 10th Dr. S.P. Ray Chaudhuri Memorial Lecture. Priorities of soil fertility and fertilizer use research in India. Journal of the Indian society of soil science. 1999; 847(4):649-660.
2. Patil YM, Sonar KR. Status of Major and Macronutrients of swell-shrink soils of Maharashtra J. Maharashtra agric. Univ. 1994; 19(2):169-172.
3. Deshmukh AH, Deshmukh PR, Harshada Changade S, Dongre VS, Solunke PS. Status of Available Boron in soils of Western Vidarbha. PKV Res. J., 2007, 31(2).
4. Jibhakate SB, Raut MM, Bhende SN, Kharche VK. Micronutrient status of soils of Katol tahasil in Nagpur District and their relationship with some soil properties. J Soils and crops. 2009; 19(1):143-146.
5. Jibhakate SB, Bhede SN, Kharche VK, Selvalakshmi V. Physico-chemical status of soils of katol Tahasil in Nagpur District, J. Soil and Crops. 2009; 19(1):122-128.
6. Ramamurthy B, Bajaj JC. Available, N, P and K status of Indian soils. Fertilizer news. 1969; 14(8):24-26.
7. Mehra RK. Effect of Application of zinc and phosphorus on the yield and Nutrient uptake (N, P, Z) by wheat (*Triticum aestivum* L.). Indian J Agric. Chem. 2003; XXX(1):1-12.
8. Katyal JC, Rattan RK. Secondary and micronutrients Research gaps and future needs. Fertilizer news. 2003; 48 (4):9-14 & 17-20 (10 pages).