



P-ISSN: 2349-8528

E-ISSN: 2321-4902

IJCS 2019; 7(6): 1180-1186

© 2019 IJCS

Received: 28-09-2019

Accepted: 30-10-2019

B Ushasri

Division of Soil Science and
Agricultural Chemistry,
Mahatma Phule Krishi
Vidyapeeth, College of
Agriculture, Kolhapur,
Maharashtra, India

DS Patil

Division of Soil Science and
Agricultural Chemistry,
Mahatma Phule Krishi
Vidyapeeth, College of
Agriculture, Kolhapur,
Maharashtra, India

RB Pawar C

Division of Soil Science and
Agricultural Chemistry,
Mahatma Phule Krishi
Vidyapeeth, College of
Agriculture, Kolhapur,
Maharashtra, India

Mukesh Kumar

Division of Soil Science and
Agricultural Chemistry,
Mahatma Phule Krishi
Vidyapeeth, College of
Agriculture, Kolhapur,
Maharashtra, India

Corresponding Author:**B Ushasri**

Division of Soil Science and
Agricultural Chemistry,
Mahatma Phule Krishi
Vidyapeeth, College of
Agriculture, Kolhapur,
Maharashtra, India

Soil fertility maps of macronutrient status of Bhudargad tehsil of Kolhapur district (M.S.)

B Ushasri, DS Patil, RB Pawar C and Mukesh Kumar

Abstract

The study was carried out to know the fertility status of major nutrients in soils of Bhudargad tehsil, Kolhapur district by using GPS-GIS technology and to correlate soil properties with the available nutrients and to evaluate the fertility index of Bhudargad tehsil. The pH of soils of Bhudargad tehsil varied from 5.5 to 8.0, most of the soils were moderately acidic (50.67%) while EC of the soils varied from 0.02 to 1.48 dS m⁻¹, and the values were normal. The calcium carbonate content of soils of Bhudargad tehsil varied from 0.50 to 2.80 per cent. Organic carbon content varied from 0.26 to 1.32 per cent and categorized as very high (6.67%), high (33.3%), moderately high (40%), moderate (13.33%) and low (6.67%). The available nitrogen, phosphorus and potassium in soils of Bhudargad tehsil ranged from 141.30 to 348.10, 7.16 to 28.00 and 100.20 to 300.00 Kg ha⁻¹, respectively. The soils of were low (57.33%) in available - nitrogen, low (17.33%) to moderately high (42.67%) in available phosphorus and moderately high (51.33%) in available potassium. The exchangeable calcium and magnesium ranged from 6.50 to 48.50 and 1.0 to 19.50 cmol (p+) Kg⁻¹, respectively. The exchangeable sodium ranged from 0.21 to 2.00 cmol (p+) Kg⁻¹. The available sulphur varied from 5.14 to 14.97 mg Kg⁻¹ respectively, and 66 per cent soil samples were in very low category.

The pH was significantly and positively correlated with potassium, calcium and magnesium. The EC was significantly and positively correlated with potassium and calcium. The organic carbon was significantly correlated with nitrogen, copper. Calcium carbonate eq. was significantly correlated with available potassium, calcium and magnesium. The fertility indices of soils of Bhudargad tehsil for organic carbon, available nitrogen, phosphorus and potassium were medium (2.10), low (1.21), low (1.62) and medium (1.94), respectively.

Keywords: Bhudargad tehsil, GPS-GIS technology, soil fertility maps

Introduction

Soil is base for every production system and knowledge on their properties, extent and specially distribution is extremely important. The soil fertility undergoes change due to intensive cropping, manuring and fertilizer applications. Application of fertilizers by the farmer in the fields without prior knowledge of soil fertility status might result in adverse effect on some soils as well as crops both in terms of nutrient deficiency and toxicity. Sing *et al.* (2005)^[17] used point estimates for districts to prepare soil fertility maps of N, P and K for Andhra Pradesh and Maharashtra states. GPS-GIS are advanced tool for studying on site specific nutrient management which can be efficiently used for monitoring soil fertility status and to prepare soil fertility maps of Bhudargad tehsil of Kolhapur district (M.S.), and it is useful for ensuring balanced fertilization to crops. Systematic study of nutrients including assessment of macro nutrient status of soil with delineation of areas of nutrient deficiency or sufficiency. The soil fertility maps are once created it is possible to transform the information about the fertility status of the area; such maps provide site specific recommendation, validation for soil fertility over the following years. Further, it is also possible to monitor the changes in fertility status of the study area with site-specific nutrient requirements of the crop. Keeping this in view, present study was undertaken in Bhudargad tehsil of Kolhapur district (M.S.).

Materials and Methods

Bhudargad tehsil of Kolhapur district was selected to assess the soil macro and micro nutrient status and delineate the fertility map, 66 villages were selected randomly in such a way that it covers the whole area of the tehsil. One hundred and fifty representative soil samples were collected from sixty six villages of Bhudargad tehsil along with GPS reading.

Collection and Processing of Soil Samples

Sixty-six villages from Bhudargad tehsil were selected for sampling keeping in mind to avoid overcrowding of sampling site on GPS based soil fertility map. Geo-referenced surface (0-22.5) cm soil samples from each selected villages representing different soils were collected. The latitude and longitude of sampling sites were recorded with the help of differential Global Positioning system with detailed observation on cropping pattern and fertilizer use. The soil samples were collected with the help of wooden peg. The samples were air dried and ground using wooden mortar and pestle and passed through 2.0 and 0.5 mm sieves. The sieved soil samples were stored in cloth bags with proper labeling for subsequent analysis. The soils were analyzed for different parameters.

The pH was measured by in 1:2.5 soil water suspension using glass electrode pH meter and EC (dS^{-1}m) was measured in the supernatant solution of 1:2.5 soil water suspension using conductivity meter (Jackson, 1973) [3]. Organic carbon by wet oxidation method (Nelson and Sommers 1982) [10]. Available Nitrogen was estimated by alkaline permanganate method of Subbiah and Asija (1956) [20] available phosphorus was extracted with 0.5 M NaHCO_3 solution buffered at pH-8.5 Watanabe and Olsen (1965) and Bray I 0.03 N ammonium fluoride at pH-3.5 (Bray and Kurtz, 1945) [1]. Available Potassium was estimated by shaking the requisite amount of soil sample with 1 N neutral ammonium Acetate solution at pH-7.0 (1:5 soil water ratio), (Knudsen and Peterson, 1982) [7]. Available and Sulphur was estimated by turbidimetry with calcium chloride extractable (Williams and Steinbergs, 1959) [23]. The CaCO_3 % by Rapid titration method (Piper, 1966) [13] and Exchangeable Ca & Mg by Versenate titration method (Page *et al.* 1982) [11].

Results and Discussion

The result of the investigation carried out during the year 2015-2016 with the view to study the GPS-GIS based fertility status of soils in Bhudargad tehsil are presented below-

Nutrient status of soils of Bhudargad tehsil

The soil samples collected from Bhudargad tehsil were analyzed by adopting standard procedure and the data pertaining to different parameters was categorized as per the six tier rating. The data pertaining to pH, Electrical conductivity, Calcium carbonate, and Organic carbon, available N, P and K are presented in Table.1 and 2.

Soil reaction (pH)

The pH of the 150 soil samples ranged from 5.50 to 8.00. Among the soil samples tested, 50.67 per cent were moderately acidic, 26 per cent were slightly acidic, 23.33 per

cent were slightly alkaline. The high rainfall and leaching of bases like Ca, Mg, Na, K was the major factor contributed to the increase in acidity in these soils.

Similar nature of observation for soil pH was also recorded by Sannappa and Manjunath (2013) [15] in soils of Western Ghats of Karnataka, India.

Electrical Conductivity (EC)

The EC ranged from 0.02 to 1.48 dS m^{-1} which indicates that most of the soils (96 per cent) were non saline in nature and are suitable for good plant growth remaining 4 per cent soil samples come under poor seed emergence category, due to accumulation of soluble salts on the soil surface and carbonates of Ca^{++} and Mg^{++} might have increased to EC more than 1 dS m^{-1} . Similar results were reported by Mandavgade *et al.* (2015) [8] for the soils of Jintur, Selu and Pathri tehsils of Parbhani district, Maharashtra.

The lower values of electrical conductivity might be due to leaching of salts from soil profile under high rainfall and undulating topographic nature.

Per cent Calcium Carbonate equivalent content

The per cent calcium carbonate equivalent content of soils ranged from 0.50-2.80 indicating barely to moderately calcareous. Majority area (61.33 per cent) of soils was slightly calcareous. In general the per cent equivalent CaCO_3 content was less than 5%, hence soils are very good for cultivation of cereals, pulses and fruit crops.

Similar nature of observation for CaCO_3 in soil series of Kolhapur district was reported by Patil (2011). The content of CaCO_3 increased from escarpment towards the foot of hill, gentle sloping of land basin and flat land topography. This might be due to leaching of soluble calcium from hill slopes towards basin area.

Organic Carbon

The organic carbon content ranged from low 0.26 (per cent) to very high (1.32 per cent) with a mean value of 0.74 per cent (moderately high). About 6.67 per cent, 13.33 per cent, 40 per cent, 33 per cent and 6.67 per cent soil samples were under low, moderate, moderately high, high and very high respectively. Similar results were reported by Jadhav (2014) [4] for Kagal Tehsil of Kolhapur district.

The low and moderate organic carbon content might be due to low input of FYM, crop residues, coarse textured soil nature and high temperatures prevailing during the summer which favours for high rate of decomposition of organic matter in soil. The mean value indicates may be moderately high organic carbon content due to cultivation of field crops, use of organic manures and fertilizers in that particular area, (Telele *et al.* 1992) [21].

Table 1: pH, EC and CaCO_3 content in soils Bhudargad tehsil

Particular	pH (1:2.5)	EC (dS m^{-1}) (1:2.5)	% CaCO_3 eq.
Mean	6.60	0.27	1.61
Range	5.50-8.00	0.02-1.48	0.5-2.8
Category	Moderately acidic 76 (50.67 %)	Normal 144 (96%)	Barely calcareous 16 (10.67%)
	Slightly acidic 39 (26%)	Poor seed emergence 6 (4 %)	Slightly calcareous 92 (61.33%)
	Slightly alkaline 35 (23.33 %)		Moderately Calcareous 42 (28%)

Total no. of soil samples-150, figures in parenthesis indicates percentage.

Available N: The available nitrogen was low in majority samples of the study area (57.33 Per cent), this might be due to high rainfall and undulating topography that leads to loss of nitates by leaching. This variation in N content might be related to soil management practices, applicaton of organic manures and fertilizers to previous crop. Similar results were reported by Pulkeshi *et al.* (2012) ^[14] in Mantagani village under transition zone of Karnataka.

Available phosphorus: The available phosphorus content was moderately high in majority of soils (42.67 Per cent) of the tehsil but it was moderate (40 Per cent) and low (17.33 Per cent) in others. The red lateritic soils showed low available phosphorus which might be due to low organic matter, higher free iron oxide content and acidic soil reaction, and formation of insoluble tri calcium phosphates under alkaline conditions. The similar trend of available phosphorus was reported by Katariya (2011) ^[6] in soils of the water management project- Block A, central campus, MPKV, Rahuri.

Available K: The available potassium in the soils ranged from 100.20 to 300.00 kg ha⁻¹ showing low to high with a mean value of 225.0 kg ha⁻¹ (moderately high). Out of soil samples collected 15.33 per cent, 6.67 per cent, 51.33 per cent and 26.67 per cent were low, moderate, moderately high and high in available K.

The available potassium content of major portion of the study area was moderately high, might be due to predominance of K rich micaceous and feldspar minerals in parent material (Pulkeshi *et al.*, 2012) ^[14]. Similar results were also recorded by Sudharani *et al.* (2013) ^[19] in rice soils of Vishakapatnam district, Andhra Pradesh.

Available Sulphur

The available sulphur is ranged from 5.14 to 14.97 mg kg⁻¹ i.e., low to moderate with a mean value of 9.01 mg kg⁻¹. About 66 per cent soil samples were low and 34 per cent samples were moderate. The low sulphur content in soils might be due to coarse textured sandy soils, leaching losses due to undulating topography and high rainfall and use of high analysis fertilizers, which are devoid of sulphur.

Table 2: Organic carbon and available nitrogen, phosphorus, potassium and sulphur content in soils of Bhudargad tehsil

Particular	Organic carbon (%)	Available nutrients (kg ha ⁻¹)			
		N	P	K	S (mg kg ⁻¹)
Mean	0.74	268.27	19.52	225.30	9.01
Range	0.26-1.32	141.30- 348.10	7.16-28.00	100.20-300.00	5.14-14.97
Very low	-	-	-	-	-
Low	10 (6.67%)	86 (57.33%)	26 (17.33%)	23 (15.33%)	99 (66 %)
Moderate	20 (13.33%)	64 (42.67%)	60 (40%)	10 (6.67%)	51 (34 %)
Moderately high	60 (40%)	-	64 (42.67%)	77 (51.33%)	-
High	50 (33.3%)	-	-	40 (26.67%)	-
Very high	10 (6.67%)	-	-	-	-

Total No. of soil samples-150, figures in parenthesis indicates percentage.

Exchangeable Calcium

The exchangeable calcium ranged from 6.50 to 48.50 [cmol(p⁺)kg⁻¹] with a mean value of 22.26 [cmol(p⁺)kg⁻¹]. Out of all the samples collected, 78.67 per cent were sufficient and 21.33 per cent samples were deficient, as the critical limit of available calcium is 15.0 [cmol(p⁺)kg⁻¹] (Durgude, 1999) ^[2].

Exchangeable Magnesium

The available magnesium ranged from 1.00 to 19.50 [cmol(p⁺)kg⁻¹] with a mean value of 6.72 [cmol(p⁺)kg⁻¹]. Out of all soil samples, 25.33 per cent were sufficient, while 74.67 per cent were deficient, as the critical limit of exchangeable magnesium is 10 [cmol(P⁺)kg⁻¹] (Durgude, 1999) ^[2]. The deficiency of magnesium might be due to the coarse textured soils and leaching losses due to heavy irrigation of some soils. Similar trend of results were observed by Mandal and Sharma (2005) ^[9] in soils of Nagpur district, (Maharashtra).




Fig 1: Location map of study area




Fig 2: pH of Bhudargad tehsil




Fig 4: CaCO₃ content of Bhudargad tehsil




Fig 3: EC of Bhudargad tehsil




Fig 5: OC of Bhudargad tehsil




Fig 6: Avail. N of Bhudargad tehsil




Fig 8: Avail. K of Bhudargad tehsil




Fig 7: Avail. P of Bhudargad tehsil




Fig 9: Avail. S of Bhudargad tehsil




Fig 10: Ca content in Bhudargad tehsil




Fig 11: Mg content in Bhudargad tehsil

Conclusion

From the study, it can be concluded that, the soils of the Bhudargad tehsil were found to be moderately acidic to slightly alkaline in reaction, normal in salt content indicating that the soils are free from salinity, low to high in organic carbon content and non-calcareous to slightly calcareous in per cent calcium carbonate equivalent content. Soils of tehsil were low to moderate in available nitrogen and that indicates nitrogen is the major limiting nutrient in the soils and also found majority samples were high (42.67%) in available phosphorus where as low (15.33%) to high (26.67%) in available potassium content. The soils were low (66%) to moderate (34%) in available sulphur and 21.33 and and 74.67 per cent soil samples found deficient in exchangeable calcium and in exchangeable magnesium content respectively. The maps generated under the study will be useful for generating homogenous units and guiding the farmers to decide the amount and kind of macronutrients to be applied for optimizing economic returns. The geo-referenced sampling sites can be revisited with the help of GPS, which helps in monitoring the changes in the status of nutrients over a period of time, which otherwise is not possible by traditional methods of sampling.

References

1. Bray RH, Kurtz LT. Determination of total, organic, and available forms of phosphorus in soils. *Soil Science*, 1945; 59:39-45.
2. Durgude AG. Morphology, Characterization, Classification and Mapping of salt affected soils of central campus, research farms, MPKV., Rahuri. Ph.D. thesis submitted to MPKV., Rahuri (M.S.), 1999.
3. Jackson ML. *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi, 1973, 256-260.
4. Jadhav RD. GPS-GIS based soil fertility maps of 'Kagal' Tehsil of Kolhapur district (M. S.) M. Sc. (Agri.) Thesis, M. P. K. V., Rahuri, 2014.
5. Kanwar JS. *In soil fertility- Theory and practical*. Indian Council of Agriculture Research, New Delhi, 1976, 202.
6. Katariya P. Characterization and classification of soils of water management project, MPKV, Rahuri. M.Sc. (Agri) Thesis, Mahatma Phule Krishi Vidyapeeth Rahuri (Maharashtra) India, 2011.
7. Knudsen DA, Peterson GA. Lithium, Sodium and Potassium, 225-246. In A. L. Page (ed). *Method soil analysis Agronomy monogram No. 9*. Am, Soc. Agron Incl. Soil Sci. Soc. Am. Publ., Madison, Wisconsin, USA, 1982.
8. Mandavgade RR, Waikar SL, Dhamak AL, Patil VD. Evaluation of micronutrient status of soils and their correlation with some chemical properties of soils of northern tehsils (Jintur, Selu and Pathri) of Parbhani district. *IOSR J Agril. and Vet. Sci.* 2015; 8(2):38-41.
9. Mandal AK, Sharma RC. Computerized data base of salt-affected soils in peninsular India using geographic information system. *J Indian Soc. Soil Sci.* 2005; 58(1):105-116.
10. Nelson DW, Sommer LE. Total carbon and organic matter. In *methods of soil analysis, Part - 2*, Page, A.L. (Ed.) Am. Soc. Agron. Inc. Soil Sci. Soc. Am. Inc. Madison, Wise. USA, 1982, 539-577.
11. Page AL. (Ed). *Methods of soil analysis Agronomy Monograph No-9. Pt.2*. American Soc. of Agron. Inc. Soil Sci Soc. of Amer. Inc. Publ. Medison, Wisconsin, USA, 1982.

12. Patil SS. Studies on soil Morphological, Chemical and Physical Properties of important soil series of Kolhapur district. M.Sc. (Agri.) Thesis, MPKV, Rahuri, 2011.
13. Piper CS. Soil and plant analysis, Hans publishers Bombay, 1966, 135-136.
14. Pulkeshi HBP, Patil PL, Dasog GS, Bidari BI, Mansur CP. Mapping of nutrient status by geographic information system in Mantagani village under northern transition zone of Karnataka. Karnataka J Agric. Sci. 2012; 25(3):332-335.
15. Sannappa B, Manjunath KG. Fertility status of soils in the selected regions of the Western Ghats of Karnataka, India. Scholars Academic Journal of Bioscience. 2013; 1(5):200-208.
16. Sharma RP, Singh M, Sharma JP. Correlation studies on micronutrients vis-à-vis soil properties in some soils of Nagaar district in semi-arid region of Rajasthan. J Indian Soc. Soil Sci. 2003; 51(4):522-527.
17. Singh KN, Raju NS, Subba Rao A, Abhishek Rathore, Sanjay Srivastava, Samanta RK *et al.* Prescribing Optimum Doses of Nutrients for Target Yield Through Soil Fertility Maps in Andhra Pradesh. J Ind. Soc. Agril. Statist. 2005; 59(2):131-140.
18. Srikanth KS, Patil PL, Dasog GS, Gali SK. Mapping of available major nutrients of a microwater -shed in northern dry zone of Karnataka. Karnataka J Agric. Sci. 2007; 21(3):391-395.
19. Sudharani Y, Jayasree G, Sesha Sai MVR. Mapping of nutrient status of rice soils in Visakhapatnam district using GIS techniques Asian Journal of Soil Science. 2013; 8:325-329.
20. Subbiah BV, Asija GL. A rapid procedure for estimation of available Nitrogen in soil. Current Science. 1956; 25:256-260.
21. Talele PE, Zende GK, Patil YM, Sonar KR. Various 103 Forms of potassium in soils of Maharashtra occurring under different Agro-climatic Zone. J potassium Res. 1992; 8(2):113-120.
22. Watanabe FS, Olsen SR. Test of ascorbic acid methods for phosphorous in water and sodium bicarbonate extract of soil. Proc. Soil Sci. Am. 1965; 21:677-678.
23. Williams, Steinbergs. Soil Sulphur fractions as chemical indices of available Sulphur in Australian soils. Australian Journal of Agriculture Research. 1959; 10:340-35.