



P-ISSN: 2349-8528

E-ISSN: 2321-4902

[www.chemijournal.com](http://www.chemijournal.com)

IJCS 2020; 8(4): 3845-3849

© 2020 IJCS

Received: 22-04-2020

Accepted: 26-06-2020

**Yedake SS**

Department of Horticulture,  
Vasantrao Naik Marathwada  
Krishi Vidyapeeth, Parbhani,  
Maharashtra, India

**Shinde SJ**

Department of Horticulture,  
Vasantrao Naik Marathwada  
Krishi Vidyapeeth, Parbhani,  
Maharashtra, India

**Waghmare PK**

Department of Agronomy,  
Vasantrao Naik Marathwada  
Krishi Vidyapeeth, Parbhani,  
Maharashtra, India

## Effect of integrated nutrient management on yield attributes and quality parameters of onion crop (*Allium cepa* L.) cv. N-53

Yedake SS, SJ Shinde, PK Waghmare and VN Shinde

DOI: <https://doi.org/10.22271/chemi.2020.v8.i4aw.10247>

### Abstract

A field experiment was carried out during the rabi season of 2018-19 at experimental farm Department of Horticulture College of Agriculture Parbhani (Maharashtra). Eleven treatments were applied with three replications and laid out under randomized block design (RBD). The result revealed that the application of 50 N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrients was recorded as best treatment for majority of characters. Therefore basis of present study it is concluded that application of Biofertilizers *Azospirillum* and *PSB* in combination with 50% compost, 50% N through chemical fertilizer, sulphur and micronutrient can be suggested cost effective combination for enhanced yield and quality onion production.

**Keywords:** nutrient management on yield and quality of onion

### Introduction

Most of the Indian population is vegetarian and vegetable plays vital role in Indian diet. India is the second largest producer of vegetables in the world next to china. Vegetables are vital sources of proteins, vitamins, phosphorus, calcium, minerals, dietary fibers, micronutrients, antioxidants and phyto-chemicals in our daily diet. Onion (*Allium cepa* L.) is one of the most important vegetable crop grown in India having both the food and medicinal values. Onion is member of *Amryllidaceae* family. It contain 2n=16 chromosome. Onion is "Queen of the kitchen" is one of the most important and commercially valuable vegetable as well as spice crop grown in India. Onion is known to check the deposition of cholesterol in blood vessels, thus protect against cure heart diseases resulting from blockage of arteries. Apart from nutrition, they also contain a wide array of potential phyto-chemicals like anti-carcinogenic principles and anti-oxidants. Onion has its own distain the world, area under onion is 4.30 million hectare with 74.26 million metric tons production in respect to 19.4 MT/ha productivity. In India, onion is grown on 1270 million hectare accounting for 21564 million tons of bulb production with 19.10 MT/ha productivity. Contribution of India in onion production in the world is around 20.2 per cent and Maharashtra is leading state in onion production in India, accounting 1064 hectare area and 218 lakh tonnes tons of total onion production and productivity of 17.9 MT/ha. (Anonymous, 2017). The Maharastra state has exported 13, 09,863.26 thousand Tonnes of fresh onion for the worth of Rs. 1,722.85 crores during year 2017 (Anonymous, APEDA Report, 2017). Therefore to increase export it is necessary to increase the productivity and quality standards of onion. The contribution of Maharashtra in onion production in India is 30.48 per cent, and specific pocket area of onion production are Nasik (Lasalgaon), Nagar (Karjat), Pune, Solapur, Satara, Dhule and Jalgaon districts on commercial scale. Nasik is the leading district for production of onion in the country.

The concept of sustainable agriculture envisages primary emphasis on manipulation and management of biological systems not only to maximize yield but also to stabilize the agro-systems and to minimize industrial input demands which may endure the adverse effect of climate change. Therefore, the experiment was conducted for sustainable production of onion with combine use of biofertilizers and chemical fertilizers. (Singh *et al.* 2017)<sup>[9]</sup>.

**Corresponding Author:****Yedake SS**

Department of Horticulture,  
Vasantrao Naik Marathwada  
Krishi Vidyapeeth, Parbhani,  
Maharashtra, India

Therefore, considering the need of time and as onion is the most important vegetable of our daily diet, it is highly essential to generate the information about the effect of different organic inorganic and biofertilization growth, yield and quality of onion. However, such type of work is scanty in onion on grown under Marathwada conditions. Hence, by keeping these points in view, the systematic investigation has been planned to evaluate the integrated nutrient management in onion. (*Allium cepa* L.)

### Material and methods

The field experiment entitled "Integrated nutrient management in Onion (*Allium cepa* L)" was carried out at Instructional-cum-Research farm of Department of Horticulture, Vasantrao Naik Marathwada Krishi Vidyapeeth, Parbhani, Maharashtra during Rabi-2018-19. The experiment consisted of eleven treatment which were laid out in randomized block design with three replication. The nursery Raised beds were prepared having size 3 x 1 x 0.15 m and upper layer of each raised bed was mixed with a mixture of well rotten and sieved FYM and sand. Raised beds were drenched with copper oxychloride. Then fungicide treated onion seeds of variety N-53 were sown on raised beds in rows at row spacing of 10 cm by mixing with fine sand. The raised beds were maintained systematically till the seedlings were ready for transplanting. Healthy, uniform 50 days old onion seedlings were transplanted at a spacing of 15 cm x 10 cm. Light irrigation was given immediately after transplanting. Compost 5 t/ha was applied before transplanting of crop to plots of all treatments. NPK were applied through urea, single super phosphate and muriate of potash by calculating as per plot size respectively. The biofertilizer were applied to onion seedlings by preparing slurry of soil with an addition of 500g of *Azospirillum* and phosphate solubilizing bacteria of each. The root of clipped onion seedling were dipped in

bioinoculant for 10 minutes and then immediately transplanted in experimental field. After sulphur and boron applied foliar spraying. Matured crop was harvested. Five plants were selected randomly from each replication to record the observations on growth and yield attributes.

Diameter of the onion bulb is the maximum distance between the opposite side of right angles to the polar diameter. It was measured by Vernier calliper in centimeter. The average length of bulb was calculated from randomly selected five fruit of each treatment with vernier calliper and value were recorded in centimetre. The polar diameter was measured from the junction of root plate to the top of the bulb from the same, which was used for recording neck thickness and mean polar diameter was worked out with the help of vernier calliper. The bulb form randomly selected five observational plants of each plot were weighed individually on an electronic balance and the average fresh weight was computed and expressed in grams. The weight of harvested and well cured bulbs from each replication and each treatment were recorded separately and average yield per plot was calculated in kilograms. The yield per hectare was calculated by multiplying the total yield per plot with hectare factor and expressed in quintals. Total acidity is determined by titrating the sample extracted in water against 0.1% sodium hydroxide. The total soluble solids content of onion bulbs was recorded with the help of hand refractometer. The average content was worked out from five observational the bulbs and it was expressed in percentage. Reducing sugars (%) and non-reducing sugars (%) of onion bulbs were determined on fresh basis as juice was taken for analysis by benedict method. The B:C ratio was calculated by using following formula.

$$B: C \text{ ratio} = \frac{\text{Gross monetary returns (Rs/ha)}}{\text{Total cost of cultivation (Rs/ha.)}}$$

**Table 1:** The treatment details

No. of Treatment	Treatments combination
T <sub>1</sub>	RDF (100:50:50 kg <sup>-1</sup> ) (control)
T <sub>2</sub>	RDF + Biofertilizers ( <i>Azospirillum</i> + <i>PSB</i> @ 500 g ha <sup>-1</sup> )
T <sub>3</sub>	RDF + Sulphur (25 kg ha <sup>-1</sup> )
T <sub>4</sub>	RDF + Micronutrients (Zinc 20 kg and Boron 5 kg ha <sup>-1</sup> )
T <sub>5</sub>	RDF + Biofertilizer + Sulphur
T <sub>6</sub>	RDF + Sulphur + Micronutrients (Zinc & Boron)
T <sub>7</sub>	RDF + Biofertilizers + Sulphur + Micronutrients
T <sub>8</sub>	Organic(100 % N- Enriched through compost 5 t ha <sup>-1</sup> + Biofertilizers)
T <sub>9</sub>	50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers
T <sub>10</sub>	50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur
T <sub>11</sub>	50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrients

## 1. Yield attributes

### 1.1 Length of bulb (cm)

The data regarding length of bulb are presented in Table 2 showed that, the significantly maximum length of bulb (6.5 cm) was observed with the treatment of T<sub>11</sub> i.e. 50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrient. It was 43.75 per cent increase as compared to control, which were statistically at par with the treatment T<sub>7</sub> (6.40cm), T<sub>10</sub> (6.20cm) and T<sub>9</sub> (6.10cm). Which was statistically followed by the treatments T<sub>4</sub> (5.40cm), T<sub>3</sub> (5.10cm), T<sub>2</sub> (5.00cm). The minimum length of bulb (3.6 cm) was observed in the treatment T<sub>1</sub> (RDF100:50:50kg ha<sup>-1</sup>) control, which was statistically at par with the treatments T<sub>8</sub> (4.10 cm), T<sub>5</sub> (4.10

cm) and T<sub>6</sub> (4.30 cm). The results were also in conformity with the findings Swati Brinjh *et al.* (2014)<sup>[7]</sup> in onion.

### 1.2 Diameter of bulb (cm)

The data regarding diameter of bulb are presented in Table 2, showed that, the significantly maximum diameter of bulb (6.60cm) was observed the treatment T<sub>11</sub> i.e. 50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrient. It was 42.43 per cent increase as compared to control, which were statistically at par with the treatments T<sub>10</sub> (6.50cm), T<sub>9</sub> (6.00cm), T<sub>7</sub> (5.60cm), T<sub>8</sub> (5.60cm) and T<sub>6</sub> (5.00cm). It was followed by the treatments T<sub>5</sub> (4.60cm), T<sub>3</sub> (4.60cm), T<sub>4</sub> (4.40cm), T<sub>2</sub> (3.80cm) While, the minimum diameter of bulb (3.90cm) was observed in the treatment T<sub>1</sub> i.e. RDF 100:50:50 kg ha<sup>-1</sup> control. The

results were also in conformity with the findings of D.N. Damse (2014)<sup>[2]</sup> in garlic.

### 1.3 Weight of bulb (g)

The data regarding weight of bulb are presented in Table 2, indicated that, maximum weight of bulb (233 g) was observed in the treatment T<sub>11</sub> (50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrient). It was 63.01 per cent increase as compared to control. Which were statistically at par with the treatments T<sub>10</sub> (213 g) and T<sub>9</sub> (211 g). It was statistically followed by with the treatments T<sub>7</sub> (158 g), T<sub>6</sub> (152 g), T<sub>8</sub> (151 g), T<sub>5</sub> (143 g), T<sub>3</sub> (114 g), T<sub>4</sub> (121 g), T<sub>2</sub> (110 g) while, the minimum (86 g) weight of bulb was observed in the treatment T<sub>1</sub> (RDF 100:50:50 kg ha<sup>-1</sup>) control. The results were also in conformity with the findings D.N. Damse (2014)<sup>[2]</sup> in garlic.

### 1.4 Yield per plot (kg)

**Table 2:** Yield attributes as influenced by integrated nutrient management in onion crop.

Tr. No	Treatments details	Bulb Length (cm)	Bulb Diameter (cm)	Weight of bulb (g)	Yield Kg/plot	Yield (q ha <sup>-1</sup> )	B:C ratio
T <sub>1</sub>	RDF(100:50:50 kg ha <sup>-1</sup> ) (control)	3.60	3.80	86	13.08	136	1.06
T <sub>2</sub>	RDF + Biofertilizers ( <i>Azospirillum</i> + <i>PSB</i> @ 500 g ha <sup>-1</sup> )	5.00	3.90	110	17.24	156	1.09
T <sub>3</sub>	RDF + Sulphur (25 kg ha <sup>-1</sup> )	5.10	4.60	114	18.42	166	1.18
T <sub>4</sub>	RDF + Micronutrients (Zinc 20 kg and Boron 5 kg ha <sup>-1</sup> )	5.40	4.40	121	20.22	183	1.40
T <sub>5</sub>	RDF + Biofertilizer + Sulphur	4.10	4.60	143	24.33	220	1.97
T <sub>6</sub>	RDF + Sulphur + Micronutrients (Zinc & Boron)	4.30	5.00	152	25.00	226	2.06
T <sub>7</sub>	RDF + Biofertilizers + Sulphur + Micronutrients	6.40	5.60	158	27.33	247	2.32
T <sub>8</sub>	Organic (100 % N- Enriched through compost 5 t ha <sup>-1</sup> + Biofertilizers)	4.00	5.60	151	26.00	235	2.88
T <sub>9</sub>	50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers	6.10	6.00	211	35.00	350	3.45
T <sub>10</sub>	50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur	6.20	6.50	213	39.66	378	3.66
T <sub>11</sub>	50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrients	6.50	6.60	233	41.66	384	3.75
	SE ±	0.35	0.61	8.04	1.60	3.58	0.069
	CD at 5%	1.04	1.81	23.77	4.75	10.59	0.206

### 1.5 Yield per hectare (q)

The data regarding yield per hectare are presented in Table 2 revealed that, the significantly highest yield per hectare (384 q) was recorded in the treatment T<sub>11</sub> i.e. 50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrient. It was 64.59 per cent increase as compared to control. Which were statistically at par with the treatment T<sub>10</sub> (378q). It was 64.03 per cent increase as compared to control. It was statistically followed by the treatments T<sub>9</sub> (350q), T<sub>7</sub> (247q), T<sub>8</sub> (235q), T<sub>6</sub> (226q), T<sub>5</sub> (220q), T<sub>4</sub> (183q), T<sub>3</sub> (166q), T<sub>2</sub> (156q) whereas, the lowest (136 q) yield per hectare was observed in the treatment T<sub>1</sub> (RDF100:50:50 kg ha<sup>-1</sup>) control. 50kg/ha<sup>-1</sup>) control.

Yield parameters highly responded to boron as well as zinc, so judicious application of zinc and boron may provide highest yield. The application of organic, inorganic and biofertilizers in various combinations. This might be due to the fact that organic manure supplied to balanced nutrition to the crop, improved soil condition and thereby resulting in better growth and development leading to higher yield. Similar results were reported by Prusty (2019)<sup>[4]</sup> in onion

## 2. Quality parameter

### 2.1 Total soluble solids (%)

The data regarding total soluble solids are presented in Table 3, indicated that the significantly highest TSS (12%) was recorded in the bulbs obtained in the treatment T<sub>11</sub> i.e. 50 % N

The data regarding yield per plot are presented in Table 2, indicated that, the significantly highest yield per plot (41.66 kg) was recorded in the treatment T<sub>11</sub> i.e. 50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrient. It was 68.61 per cent increase as compared to control. Which were statistically at par with the treatment T<sub>10</sub> (39.66 kg). It was 67.02 per cent increase as compared to control. Which was statistically followed by the treatments T<sub>9</sub> (35.00 kg), T<sub>7</sub> (27.33 kg), T<sub>8</sub> (26.00 kg), T<sub>6</sub> (25.00 kg), T<sub>5</sub> (24.33 kg), T<sub>4</sub> (20.22 kg), T<sub>3</sub> (18.42 kg), T<sub>2</sub> (17.24 kg) While, the lowest yield (13.08 kg) per plot was observed in the treatment T<sub>1</sub> (RDF 100:50:50 kg ha<sup>-1</sup>) control. Which was statistically at par with the treatment T<sub>2</sub> (17.24 kg). Similar results were reported by Prusty (2019)<sup>[4]</sup> in onion.

Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrient, which were statistically at par treatments T<sub>10</sub> (11.30%), T<sub>9</sub> (11.30%) and T<sub>8</sub> (11.00%). Which was statistically followed by with treatments T<sub>7</sub> (10%), T<sub>6</sub> (9.60%), while the minimum TSS (8%) was observed in the treatment T<sub>1</sub> (RDF100:50:50 kg ha<sup>-1</sup>) control treatment. which were statistically at par with the treatment T<sub>3</sub> (8.10 %), T<sub>5</sub> (8.30 %), T<sub>2</sub> (8.40 %), T<sub>4</sub> (8, 60 %), and T<sub>6</sub> (9.60 %). Combination of biofertilizers, organic and inorganic sources of nutrient found beneficial in increasing total soluble solids. This may be due to the known fact that, organic and inorganic nutrient sources are capable for supplying adequate macro and micro nutrients which plays major role in quality improvement through desirable enzymatic changes took place during entire plant growth. The results are in consequences with the findings of, S. Ghanti *et al.* (2009)<sup>[6]</sup>.

### 2.2 Acidity (%)

The data regarding acidity are presented in Table 13, indicated that, the significantly highest acidity (2.63%) was recorded in the bulbs obtained in the treatment T<sub>11</sub> i.e. 50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrient, which were statistically at par with the treatments T<sub>5</sub> (2.50%), T<sub>9</sub> (2.43%), T<sub>4</sub> (2.40%) and T<sub>10</sub> (2.36%). The minimum acidity (1.6%) was observed in the treatment T<sub>1</sub> (RDF100:50:50 kg ha<sup>-1</sup>) control.

which were statistically at par with the treatments T<sub>2</sub> (1.76 %), T<sub>8</sub> (1.70 %), T<sub>7</sub> (1.90 %) and T<sub>6</sub> (2.00 %). Addition of biofertilizers, sulphur and micronutrients along with general recommended dose of fertilizers. Has significant effect on onion acidity. The results are in consequences with the findings Heerendra Prasad *et al.* (2017)<sup>[3]</sup> in onion.

### 2.3 Total sugars (%)

The data regarding total sugar percentage are presented in Table 14, indicated that, maximum total sugars (9.07%) was recorded in the onion bulbs produced in the treatment T<sub>11</sub> i.e. 50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrient), which were statistically at par with the treatment T<sub>10</sub> (9.05%), T<sub>8</sub> (8.98%) T<sub>7</sub> (8.91%), T<sub>3</sub> (8.75%), T<sub>6</sub> (8.69%), T<sub>4</sub> (8.69%), T<sub>9</sub> (8.52%) and T<sub>5</sub> (7.81%). The minimum total sugars (7.11%) were observed in the treatment T<sub>1</sub> (RDF100:50:50 kg ha<sup>-1</sup>) control, which were statistically at par with the treatment T<sub>2</sub> (7.76 %).

### 2.4 Reducing sugars (%)

The data regarding reducing sugar percentage are presented in Table 14, indicated that, maximum reducing sugars (3.93%) was recorded in the onion bulbs produced in the treatment T<sub>11</sub> i.e. 50 % N Enriched through compost + Biofertilizers + 50 %

N through chemical fertilizers + Sulphur + Micronutrient, which was statistically at par with T<sub>10</sub> (3.91%), T<sub>9</sub> (3.88%), T<sub>8</sub>(3.86%), T<sub>7</sub>(3.81%), T<sub>5</sub> (3.76%), T<sub>3</sub> (3.75 %), T<sub>4</sub> (3.76%) and T<sub>6</sub>(3.76%). The minimum (3.28%) reducing sugars was observed in the treatment T<sub>1</sub> (RDF 100:50:50 kg ha<sup>-1</sup>) control treatment. which were statistically at par with the treatment T<sub>2</sub> (3.32 %).

### 2.5 Non reducing sugar (%)

The data regarding non reducing sugar percentage are presented in Table 14 indicated that the significantly maximum non reducing sugars (5.15%) was recorded in the onion bulbs produced in the treatment T<sub>11</sub> i.e. 50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrient, which were statistically at par with the treatments T<sub>10</sub>(5.13%), T<sub>7</sub>(5.11 %), T<sub>8</sub>(5.10 %), T<sub>3</sub> (5.03 %), T<sub>6</sub>(5.00 %), T<sub>4</sub> (5.00 %) and T<sub>9</sub> (4.80 %) While, the minimum non reducing sugars (3.91%) were observed in the treatment T<sub>1</sub> (RDF 100:50:50 kg ha<sup>-1</sup>) control. The application of biofertilizers, inorganic and organic manure improved the quality parameters in respect of TSS, total sugar, reducing sugar and non-reducing sugar content. The results are in consequences with the findings S. Ghanti *et al.* (2009)<sup>[6]</sup>, and Anil kumar *et al.* (2017)<sup>[1]</sup> in onion.

**Table 3:** Quality parameters as influenced by integrated nutrient management in onion crop

Tr. No	Treatments details	TSS (%)	Acidity (%)	Reducing sugar (%)	Nonreducing sugar (%)	Total sugar (%)
T <sub>1</sub>	RDF (100:50:50 kg ha <sup>-1</sup> ) (control)	8.0	1.60	3.28	3.91	7.11
T <sub>2</sub>	RDF + Biofertilizers ( <i>Azospirillum</i> + <i>PSB</i> @ 500 gha <sup>-1</sup> )	8.4	1.76	3.32	4.04	7.36
T <sub>3</sub>	RDF + Sulphur (25 kg ha <sup>-1</sup> )	8.1	2.03	3.75	5.04	8.75
T <sub>4</sub>	RDF + Micronutrients (Zinc 20 kg and Boron 5 kg ha <sup>-1</sup> )	8.6	2.40	3.69	5.00	8.69
T <sub>5</sub>	RDF + Biofertilizer + Sulphur	8.3	2.50	3.76	4.06	7.81
T <sub>6</sub>	RDF + Sulphur + Micronutrients (Zinc & Boron)	9.6	2.00	3.69	5.00	8.69
T <sub>7</sub>	RDF + Biofertilizers + Sulphur + Micronutrients	10.0	1.90	3.81	5.11	8.91
T <sub>8</sub>	Organic (100 % N- Enriched through compost 5 t ha <sup>-1</sup> + Biodfertilizers)	11.0	1.70	3.86	5.10	8.98
T <sub>9</sub>	50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers	11.3	2.43	3.88	4.80	8.52
T <sub>10</sub>	50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur	11.3	2.36	3.91	5.13	9.05
T <sub>11</sub>	50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrients	12	2.63	3.93	5.15	9.07
	SE ±	0.59	0.14	0.13	0.24	0.44
	CD at 5%	1.77	0.41	0.39	0.72	1.32

### Conclusion

The overall assessment of the results of present investigation entitled "Integrated nutrient management in onion (*Allium cepa* L.) Cv. N-53" concluded that the treatment T<sub>11</sub> (50 % N Enriched through compost + Biofertilizers + 50 % N through chemical fertilizers + Sulphur + Micronutrients) significantly influenced the yield and quality attributes and benefit cost ratio of onion. The growth parameter *viz.* bulb parameter like length of bulb, diameter of bulb, weight of bulb, yield parameter *viz.* yield per plot and yield per hectare, quality parameter *viz.* total soluble solids, acidity, total sugar, reducing sugar and non-reducing sugar.

### References

- Anil Kumar, Ram RB, Sutanumazi. Effect of organic manures, biofertilizers and micronutrients on growth, yield and quality of onion (*Allium cepa* L.) International Journal of Agricultural Sciences. 2017; 13(2):236-241.
- Damse DN, Bhalekar MN, Pawar PK. Effect of integrated nutrient management on growth and yield of garlic. The bioscan.2014; 9(4):1557-1560.
- Heerendra Prasad, Paramjeet Sajwan, Meena Kumari, SPS Solanki. Effect of Organic Manures and Biofertilizer on Plant Growth, Yield and Quality of Horticultural Crop: A Review- International Journal of Chemical Studies.2017; 5(1):217-221.
- Prusty M, Mishra N, Kar DS, Pal S. Effect of integrated nutrient management on growth and yield of onion (*Allium cepa* L.) cv Bhima super International Journal of Agriculture Sciences. 2019; 11(4):7910-7912.
- Ram Vachan, Tripathi SM. Study on the effect of bio-fertilizer with chemical fertilizer on plant growth, Yield and economics of Rabi season onion (*Allium cepa* L.) cv. NHRDFRed2. Journal of Pharmacognosy and Phytochemistry. 2017; 6(5):1496-1499.
- Ghanti S, Sharangi AB. Effect of bio-fertilizers on growth, yield and quality of onion cv. Sukhsagar Journal of Crop and Weed.2009; 5(1):120-123.

7. Swati Brinjh, Sanjay Kumar, Devendra Kumar, Manoj Kuamar. Effect of integrated nutrient management on growth yield and quality in onion cv. Pusamadhavi Plant Archives. 2014; 14(1):557-559.
8. Umesh Acharya, Narayan Sharma Rimal, Venkatesan K, Saraswathi T, Subramanian KS. Response on growth, yield and quality parameters of Multiplier Onion (*Allium cepa* L. varaggregatum Don.) var. CO (On) 5 with different doses and method of Zinc and Boron application. International Journal of Agronomy and Agricultural Research. 2015; 6(4):59-67.
9. Singh VK, Amrita Kumari, Chaudhary VK, Shree S. Role of Biofertilizer and Chemical Fertilizer for Sustainable Onion (*Allium cepa* L.) Production. International Journal of Current Microbiology and Applied Sciences. 2017; 6:2034-2040.